

Species	Date	River Section	Number	Approximate Size (inches)
Bonytail	Oct 1998	Colorado	3,280	5
	Oct 1998	Lower Green	3,000	5
	Mar 1999	Colorado	15 (with radio tags)	10
	Apr 1999	Colorado	10,000	4
	Apr 1999	Lower Green	10,000	4
	Mar 2000	Lower Green	13	
	Apr 2000	Lower Green	19,987	4-7
	Apr 2000	Colorado	15,037	3
	Jul 2000	Yampa	5,000	4
	Jul 2000	Middle Green¹	5,000	4
Razorback sucker	Oct 1998	Middle Green ²	125	6-8
	Apr & Aug 1999	Middle Green ²	6659	4-8
	May 1999	Middle Green ²	57,900	<1
	May 1999	Middle Green	35 (with radio tags)	>10
	Jun 1999	Middle Green	738	10-16
	Jun 2000	Old Charlie	9599	<1
	Jun 2000	Middle Green	79	17
	Jun 2000	Stewart Lake	145	12
	Jun 2000	Old Charlie	2,106	>6
	Sept 1998	Gunnison	249	9
	Oct 1998	Gunnison	126	16
	May & Nov 1999	Gunnison	2,772	8
	Sept-Oct 1999	Colorado ³	3,498	8
	Apr 2000	Colorado³	7,147	4-6
	Aug 2000	Colorado³	3,875	4-13
	Aug 2000	Gunnison	1,640	3-13
Colorado pikeminnow	Apr 1999	Middle Green	36	16
	Jun 2000	Colorado	60⁴	17-22

¹ This reach of the middle Green River was in the Lodore Canyon.

² These smaller fish were stocked in depression wetlands where early life stages can take advantage of resources for growth and protection.

³ Colorado River at Parachute, CO.

⁴ These fish are 1991 year class from Horsethief ponds and are part of the translocation study